



ESO

# Riesen-Schmiede

## Die Entstehung der massereichsten Sterne

Von Henrik Beuther

### Weiterführende bibliografische Angaben

- [9] Yorke, H., Sonnhalter, C.: On the formation of massive stars. In: *Astrophysical Journal* 569, S. 846–862, 2002.
- [10] Krumholz, M., Klein, R., McKee, C.: Radiation-hydrodynamic simulations of collapse and fragmentation in massive protostellar cores. In: *Astrophysical Journal* 656, S. 959–979, 2007.
- [11] Bonnell, I., Larson, R., Zinnecker, H.: The origin of the initial mass function. In: *Protostars and Planets V*, S. 149–164, 2007.
- [12] Bally, J., Zinnecker, H.: The birth of high-mass stars: Accretion and/or mergers? In: *Astronomical Journal* 129, S. 2281–2293, 2005.
- [13] Zinnecker, H. und Yorke H.: Toward understanding massive star formation. In: *Annual Review of Astronomy and Astrophysics* 45, S. 481–563, 2007.
- [14] Krumholz, M. und McKee, C.: A minimum column density of  $1 \text{ g cm}^{-2}$  for massive star formation. In: *Nature* 451, S. 1082–1084, 2008.
- [15] Fallscheer C., Beuther H., Zhang Q. und Sridharan T.K.: Rotational structure and outflow in the infrared dark cloud 18223-3. In: *Astronomy and Astrophysics* 504, S. 127–137, 2009.
- [16] Rodon, J., Beuther, H., Schilke, P.: Fragmentation in the massive star-forming region IRAS 19410: A core mass function resembling the IMF, eingereicht bei *Astronomy and Astrophysics*.
- [17] Beuther, H., Zhang, Q., Greenhill, L. et al.: Line imaging of Orion-KL at  $865 \mu\text{m}$  with the submillimeter array. In: *Astrophysical Journal* 632, S. 355–370, 2005.
- [18] Krumholz, M., Klein, R., McKee, C.: Molecular line emission from massive protostellar disks: Predictions for ALMA and EVLA. In: *Astrophysical Journal* 665, S. 478–491, 2007.
- [19] Beuther, H., Churchwell, E., McKee, C., Tan J.: The formation of massive stars. In: *Protostars and Planets V*, S. 165–180, 2007.